

Integro-differential equations linked to compound birth processes with infinitely divisible addends

Luisa Beghin* Janusz Gajda† Aditya Maheshwari‡

Abstract

Stochastic modelling of fatigue (and other material's deterioration), as well as of cumulative damage in risk theory, are often based on compound sums of independent random variables, where the number of addends is represented by an independent counting process. We consider here a cumulative model where, instead of a renewal process (as in the Poisson case), a linear birth (or Yule) process is used. This corresponds to the assumption that the frequency of “damage” increments accelerates according to the increasing number of “damages”. We start from the partial differential equation satisfied by its transition density, in the case of exponentially distributed addends, and then we generalize it by introducing a space-derivative of convolution type (i.e. defined in terms of the Laplace exponent of a subordinator). Then we are concerned with the solution of integro-differential equations, which, in particular cases, reduce to fractional ones. Correspondingly, we analyze the related cumulative jump processes under a general infinitely divisible distribution of the (positive) jumps. Some special cases (such as the stable, tempered stable, gamma and Poisson) are presented.

Keywords: Integro-differential equations, Convolution-type derivatives, Cumulative damage models, First-passage time, Infinitely divisible laws.
AMS Mathematical Subject Classification (2010). 35R09, 35R11, 60G50, 33E12, 26A33.

1 Introduction

Compound counting processes, especially in the Poisson case, are widely studied and applied in many different fields: in reliability theory (for studying the development of fatigue in materials) and in collective risk theory, where many cumulative damage models are defined in terms of the following sum

$$Y(t) := \sum_{j=0}^{N(t)} X_j. \quad (1.1)$$

The addends X_j are assumed to be i.i.d. random variables, for $j = 1, 2, \dots$, and $N := \{N(t), t \geq 0\}$ is an independent counting (i.e. a non-negative, integer valued and non-decreasing) stochastic process. In the special case where N is a homogeneous Poisson process, with rate λ , the well-known compound Poisson process is obtained.

*Sapienza University of Rome, P.le A. Moro 5, 00185 Roma, Italy. e-mail: luisa.beghin@uniroma1.it

†University of Warsaw, Poland

‡Indian Institute of Management Indore, India

Under the assumption that the counting process N is Poisson with parameter $\lambda > 0$, and the jumps are exponentially distributed with parameter ξ , it has been proved in [2] that the distribution of the compound Poisson process defined in (1.1), can be written in terms of a Wright function. Moreover the density of its absolutely continuous component $f_Y(y, t)$, for $y, t > 0$ satisfy the following differential equation

$$\xi \frac{\partial}{\partial t} f = - \left[\lambda + \frac{\partial}{\partial t} \right] \frac{\partial}{\partial y} f, \quad (1.2)$$

with conditions

$$\begin{cases} f(y, 0) = 0 \\ \int_0^{+\infty} f(y, t) dy = 1 - e^{-\lambda t} \end{cases} \cdot \quad (1.3)$$

This process has been generalized in [2] to the fractional case and in [3] the assumption of Gamma distributed jumps has been considered.

We consider here the cumulative process (1.1), when the number of addends, instead of being represented by a renewal counting process (as in the Poisson case), is assumed to be a linear birth (or Yule) process $B := \{B(t), t \geq 0\}$, with one progenitor and with rate $\lambda_k = k\lambda$, for $k = 1, 2, \dots$, i.e.

$$Y(t) = \sum_{j=1}^{B(t)} X_j, \quad t \geq 0, \quad (1.4)$$

under the assumption that X_j are i.i.d. positive random variables, independent from B . This model can be also described by assuming that each member of a population gives birth independently to one offspring at an exponential time with rate λ . Moreover each member of the population produces a random (positive) “damage” that contribute individually to the “total damage” of the population. Alternatively, we can assume that X_j ’s represent the claim size of the j -th policy holder and that the number of claims for the insurance company evolves in time according to a birth process $B(t)$, i.e. when, for example, the arrival rate of the new claims is proportional to the number of the claims previously arrived. In this case a crucial random variable is represented by the time to ruin of the company, i.e.

$$\tau := \inf\{t : Y(t) > ct + u\}, \quad u, c > 0,$$

where u is the initial capital and c is constant risk premium rate. For applications to the risk theory of compound birth processes, see the very recent paper [22].

This kind of process has been introduced by [23] for modelling crack growth with accelerating frequency of increments as the crack grows. Also in the cumulative models applied to reliability theory, the study of the first-passage time through a certain critical value, i.e. $T_\beta = \inf\{t \geq 0 : Y(t) > \beta\}$, is crucial. In particular, T_β is the fatigue failure time, in the cumulative fatigue model. Since, in case of non-negative addends, $Y(t)$ is non-decreasing for any t , we get

$$P\{T_\beta > t\} = P\{Y(t) < \beta\} = F_Y(\beta, t). \quad (1.5)$$

We recall that the birth process B is a continuous-time Markov process and, in the linear case, its probability mass function is given by

$$p_n(t) := P \{B(t) = n | B(0) = 1\} = e^{-\lambda t} \left(1 - e^{-\lambda t}\right)^{n-1}, \quad t \geq 0, n = 1, 2, \dots$$

In the fractional case, the birth processes have been studied in [19], [20] and [4]. In [14] a non-markovian generalization of the Yule process has been introduced.

We start by deriving the partial differential equation satisfied by the transition density of (1.4) in the case of exponentially distributed addends; then we generalize it by introducing a space-derivative \mathcal{D}_y^g of convolution-type (defined by means of the Laplace exponent $g(\theta)$ of a subordinator). More precisely, we will be concerned with the the solution to the integro-differential equation

$$\frac{\partial}{\partial t} f(y, t) = -\lambda \frac{e^{\lambda t}}{\xi} \mathcal{D}_y^g [f * f](y, t), \quad y, t, \lambda, \xi > 0, \quad (1.6)$$

under certain initial conditions, where we denote by $f_1 * f_2$ the convolution of the functions f_1 and f_2 . For the (integral) definition of \mathcal{D}_y^g see (1.8) below.

We will prove that the solution of (1.6) coincides with the density f_{Y_g} of (1.4) when the distribution of the addends is extended (from the exponential case) inside the class of infinitely divisible laws. Note that, for $g(\theta) = \theta^\alpha$ and $\alpha \in (0, 1)$, the derivative \mathcal{D}_y^g reduces to a Caputo fractional derivative of order α (see (1.8) below) and then equation (1.6) becomes a fractional differential equation. Some special cases (such as the stable, tempered stable, gamma and Poisson cases) will be illustrated. As we will see, our model will prove to be more flexible and adaptable to the real data provided that the appropriate distributions of the addends and the corresponding parameters' values are chosen.

Let $g : (0, +\infty) \rightarrow \mathbb{R}$ be a Bernstein function, i.e. let g be non-negative, infinitely differentiable and such that, for any $x \in (0, +\infty)$,

$$(-1)^n \frac{d^n}{dx^n} g(x) \leq 0, \quad \text{for any } n \in \mathbb{N}.$$

A function g is a Bernstein function if and only if it admits the following representation

$$g(x) = a + bx + \int_0^{+\infty} (1 - e^{-sx}) \bar{\nu}(ds),$$

for $a, b \in \mathbb{R}$, where $\bar{\nu}$ is the corresponding Lévy measure and $(a, b, \bar{\nu})$ is called the Lévy triplet of g . Then a subordinator is the stochastic process with non-decreasing paths $\mathcal{A}_g := \{\mathcal{A}_g(t), t \geq 0\}$, such that

$$\mathbb{E} e^{-\theta \mathcal{A}_g(t)} = e^{-g(\theta)t}, \quad (1.7)$$

i.e. $g(\theta)$ is the Laplace exponent of \mathcal{A}_g . Let moreover $\mathcal{L}_g(t), t \geq 0$, be its inverse, i.e.

$$\mathcal{L}_g(t) = \inf \{s \geq 0 : \mathcal{A}_g(s) > t\}, \quad t > 0$$

and $l_g(x, t) = \Pr \{\mathcal{L}_g(t) \in dx\} / dx$ be its transition density.

We recall the definition of the convolution-type derivative on the positive half-axes, in the sense of Caputo (see [24], Def.2.4, for $b = 0$) :

$$\mathcal{D}_t^g u(t) := \int_0^t \frac{d}{ds} u(t-s) \nu(s) ds, \quad t > 0, \quad (1.8)$$

where ν is the tail of the Lévy measure $\bar{\nu}$, i.e. $\nu(s) = \int_s^{+\infty} \bar{\nu}(dz)$. Convolution-type derivatives (or derivatives defined as integrals with memory kernels) have been treated recently by many authors: see, among the others, [13], [8], [21].

The Laplace transform of \mathcal{D}_t^g is given by

$$\int_0^{+\infty} e^{-\theta t} \mathcal{D}_t^g u(t) dt = g(\theta) \tilde{u}(\theta) - \frac{g(\theta)}{\theta} u(0), \quad \mathcal{R}(\theta) > \theta_0, \quad (1.9)$$

(see [24], Lemma 2.5). It is easy to check that, in the trivial case where $g(\theta) = \theta$, the convolution-type derivative coincides with the first-order derivative, while, for $g(\theta) = \theta^\alpha$, for $\alpha \in (0, 1)$, it coincides with the Caputo fractional derivative (see e.g. [12], p.90) of order α .

2 The exponential case

As a preliminary result, we consider the case of the compound birth process Y with exponentially distributed addends: see [6] and [23] for possible applications, in survival analysis and reliability theory, respectively.

Lemma 2.1 *Let X_j be i.i.d. $\text{Exp}(\xi)$, for $j = 1, 2, \dots$, then the density of Y , defined in (1.4), i.e.*

$$f_Y(y, t) = \xi \exp \left\{ -\lambda t - \xi e^{-\lambda t} y \right\}, \quad y, t \geq 0, \quad (2.1)$$

satisfies the equation

$$\frac{\partial}{\partial t} f = -\lambda \frac{\partial}{\partial y} (y f), \quad y, t \geq 0, \quad (2.2)$$

with $f(y, 0) = f_X(y) = \xi e^{-\xi y}$.

Proof. By a conditioning argument, we can write

$$f_Y(y, t) = \sum_{n=1}^{\infty} p_n(t) f_X^{*(n)}(y),$$

which, under Laplace transform, gives

$$\begin{aligned} \tilde{f}_Y(\theta, t) &= \sum_{n=1}^{\infty} p_n(t) \tilde{f}_X^{*(n)}(\theta) = \sum_{n=1}^{\infty} p_n(t) \left[\tilde{f}_X(\theta) \right]^n \\ &= e^{-\lambda t} \sum_{n=1}^{\infty} \left(1 - e^{-\lambda t} \right)^{n-1} \left[\frac{\xi}{\xi + \theta} \right]^n = \frac{\xi e^{-\lambda t}}{\theta + \xi e^{-\lambda t}}. \end{aligned} \quad (2.3)$$

It can be easily checked that (2.1) satisfies equation (2.2) and the initial condition. ■

In Fig.1 we plot the probability density function (hereafter pdf) of the process Y , defined in (1.4) (in the case of exponentially distributed addends), estimated directly from the realizations of the process. Moreover we compare it with the theoretical pdf given in (2.1), for different values of t . One can notice the perfect agreement between the theoretical and empirical pdf's.




Figure 1: Compound birth process Y , with exponential distributed addends: theoretical and empirical pdf's for different values of t and for $\lambda = \xi = 1$.

Remark 2.1 Equation (2.2) can be considered as special case of the Fokker-Planck equation with null diffusion coefficient. Moreover, it can be alternatively written as

$$\frac{\partial}{\partial t} f = -\lambda \frac{e^{\lambda t}}{\xi} \frac{\partial}{\partial y} (f * f), \quad y, t \geq 0. \quad (2.4)$$

Indeed, we have that

$$\begin{aligned} \frac{e^{\lambda t}}{\xi} \frac{\partial}{\partial y} (f_Y * f_Y) &= \frac{e^{\lambda t}}{\xi} \xi^2 e^{-2\lambda t} \frac{\partial}{\partial y} \int_0^y \exp\{-\xi e^{-\lambda t} z\} \exp\{-\xi e^{-\lambda t} (y-z)\} dz \\ &= \xi e^{-\lambda t} \frac{\partial}{\partial y} (y \exp\{-\xi e^{-\lambda t} y\}) = \frac{\partial}{\partial y} (y f_Y). \end{aligned}$$

In the next section, we will generalize the governing equation of our model in the form given in (2.4).

The distribution of the first-passage time of Y through the level $\beta > 1$ can be written as follows:

$$P\{T_\beta < t\} = \begin{cases} 0, & t \leq 0 \\ \exp\{-\xi e^{-\lambda t} \beta\}, & t > 0 \end{cases} \quad (2.5)$$

by considering (1.5). The jump in zero of (2.5) is equal to $P\{T_\beta = 0\} = P\{X > \beta\} = e^{-\xi\beta}$. The absolutely continuous component thus displays a Gumbel-type distribution. Its mean

value can be obtained as follows:

$$\begin{aligned}
\mathbb{E}T_\beta &= \int_0^{+\infty} \left[1 - \exp \left\{ -\xi e^{-\lambda t} \beta \right\} \right] dt = - \sum_{j=1}^{\infty} \frac{(-\xi\beta)^j}{j!} \int_0^{+\infty} e^{-\lambda t j} dt \quad (2.6) \\
&= -\frac{1}{\lambda} \sum_{j=1}^{\infty} \frac{(-\xi\beta)^j}{j^2(j-1)!} = \frac{\xi\beta}{\lambda} \sum_{l=0}^{\infty} \frac{(-\xi\beta)^l}{l!} \frac{\Gamma(l+1)^2}{\Gamma(l+2)^2} \\
&= \frac{\xi\beta}{\lambda} {}_2\Psi_2 \left[-\xi\beta \mid \begin{matrix} (1,1) & (1,1) \\ (2,1) & (2,1) \end{matrix} \right]
\end{aligned}$$

where

$${}_p\Psi_q \left[x \mid \begin{matrix} (a_l, \alpha_l)_{1,p} \\ (b_l, \beta_l)_{1,q} \end{matrix} \right], \quad x, a_l, b_j \in \mathbb{C}, \alpha_l, \beta_j \in \mathbb{R}, l = 1, \dots, p, j = 1, \dots, q$$

for $p, q \in \mathbb{N}$, is the Fox-Wright function (see [12], p.56). The asymptotic behavior of $\mathbb{E}T_\beta$ can be studied by considering formula (1.12.68), p.67 in [12], so that we can write

$$\mathbb{E}T_\beta = \frac{\xi\beta}{\lambda} H_{2,3}^{1,2} \left[\xi\beta \mid \begin{matrix} (0,1) & (0,1) \\ (0,1) & (-1,1) & (-1,1) \end{matrix} \right],$$

where

$$H_{p,q}^{m,n} \left[x \mid \begin{matrix} (a_p, A_p) \\ (b_q, B_q) \end{matrix} \right], \quad x, a_i, b_j \in \mathbb{C}, A_i, B_j \in \mathbb{R}^+, i = 1, \dots, p, j = 1, \dots, q$$

for $p, q, m, n \in \mathbb{N}$, with $0 \leq n \leq p, 1 \leq m \leq q$, is the H-function (see e.g. [17], p.2). Then, by applying Theorem 1.2, p.19 in [17], for $\mu = \alpha = 1 > 0$ and $d = \min \{-1, -1\} = -1$ we get, for $\beta \rightarrow +\infty$,

$$\mathbb{E}T_\beta \simeq \frac{\xi\beta}{\lambda} O(\beta^{-1}) = \frac{\xi}{\lambda} O(1).$$

3 The infinitely divisible cases

We now extend the previous results, by considering the relaxation equation with the convolution-type derivative defined in (1.8). Indeed, it is well-known that the survival (or reliability) function $\Phi_X(x) := \int_x^{+\infty} f_X(u) du$ of the r.v. $X \sim \text{Exp}(\xi)$ satisfies the so-called relaxation equation

$$\frac{d}{dx} u(x) = -\xi u(x), \quad x \geq 0, \quad (3.1)$$

with initial condition $u(0) = 1$. As we will see later, replacing the space-derivative with \mathcal{D}_x^g corresponds to generalize the distribution of X , inside the class of infinitely divisible r.v.'s.

Lemma 3.1 *Let \mathcal{A}_g denote the subordinator defined by (1.7), then the initial-value problem*

$$\begin{cases} \mathcal{D}_x^g u(x) = -\xi u(x) \\ u(0) = 1, \end{cases} \quad (3.2)$$

with $x \geq 0, \xi > 0$, is satisfied by

$$u(x) = \xi \int_0^{+\infty} e^{-\xi t} P \{ \mathcal{A}_g(t) \geq x \} dt. \quad (3.3)$$

Proof. We take the Laplace transform of (3.2), by considering (1.9), and get, for $\mathcal{R}(\theta) > \theta_0$,

$$g(\theta)\tilde{u}(\theta) - \frac{g(\theta)}{\theta}u(0) = -\xi\tilde{u}(\theta),$$

so that

$$\tilde{u}(\theta) = \frac{g(\theta)}{\theta} \frac{1}{g(\theta) + \xi}. \quad (3.4)$$

On the other hand, from (3.3) we get

$$\begin{aligned} \tilde{u}(\theta) &= \xi \int_0^{+\infty} e^{-\theta x} dx \int_0^{+\infty} e^{-\xi t} dt \int_x^{+\infty} h_g(y, t) dy = \xi \int_0^{+\infty} e^{-\xi t} dt \int_0^{+\infty} h_g(y, t) dy \int_0^y e^{-\theta x} dx \\ &= \frac{\xi}{\theta} \int_0^{+\infty} e^{-\xi t} dt \int_0^{+\infty} h_g(y, t) (1 - e^{-\theta y}) dy = \frac{\xi}{\theta} \left[\frac{1}{\xi} - \frac{1}{\xi + g(\theta)} \right], \end{aligned}$$

which is equal to (3.4). The initial condition is clearly satisfied by (3.3). ■

In the trivial case $g(\theta) = \theta$, the differential equation in (3.2) reduces to (3.1), which is satisfied by $\Phi_X(x) = e^{-\xi x}$. For $g(\theta) = \theta^\alpha$, $\alpha \in (0, 1)$, equation (3.2) coincides with the well-known fractional relaxation equation (see e.g. [9], [16] and [1]). In all the other cases, the solution corresponds to the survival function of a subordinator stopped at an independent exponential time, i.e. $\mathcal{A}_g(X)$.

Alternatively, (3.3) gives the probability that the subordinator \mathcal{A}_g hits (or crosses) a certain level in a random time smaller than an exponentially distributed one. Indeed, by considering that

$$P\{\mathcal{A}_g(t) \geq x\} = P\{\mathcal{L}_g(x) \leq t\},$$

we can write the solution of (3.2) as

$$u(x) = \xi \int_0^{+\infty} e^{-\xi t} P\{\mathcal{L}_g(x) \leq t\} dt = P\{\mathcal{L}_g(x) \leq X\}, \quad (3.5)$$

with $X \sim \text{Exp}(\xi)$. In the special case where $g(\theta) = \sqrt{\theta}$, the inverse stable subordinator \mathcal{L}_g is equal in distribution to a Brownian motion reflecting in the origin and thus (3.5) reduces to the probability that the Brownian motion is under an exponentially distributed barrier at time x , i.e, $P\{|B(x)| \leq X\}$ (see [1]).

Let us now consider the r.v.'s X_j^g , for $j = 1, 2, \dots$, with survival function $\Phi_{X^{(g)}}$ coinciding with (3.3). They are thus non-negative, infinitely divisible, with the following Laplace exponent

$$\psi_{X^{(g)}}(\theta) := -\log \mathbb{E}e^{-\theta X^{(g)}} = \log \left(1 + \frac{g(\theta)}{\xi} \right), \quad \xi, \theta > 0. \quad (3.6)$$

Clearly, in the special case $g(\theta) = \theta$, formula (3.6) reduces to the Laplace exponent of the exponential law. Thus we assume here that the density of the addends in (1.1) is given by

$$f_{X^{(g)}}(x) = \xi \int_0^{+\infty} e^{-\xi t} h_g(x, t) dt, \quad \xi, y > 0, \quad (3.7)$$

(where $h_g(\cdot, t)$ is the density of the subordinator \mathcal{A}_g), or, alternatively, that the following equality in law holds : $X^{(g)} \stackrel{d}{=} \mathcal{A}_g(X)$.

Theorem 3.2 *The solution to equation*

$$\frac{\partial}{\partial t} f(y, t) = -\lambda \frac{e^{\lambda t}}{\xi} \mathcal{D}_y^g [f * f](y, t), \quad y, t > 0, \quad (3.8)$$

under the initial condition $f(y, 0) = f_{X^{(g)}}(y)$, is given by the density function of the process

$$Y_g(t) = \sum_{j=1}^{B(t)} X_j^{(g)}, \quad t \geq 0. \quad (3.9)$$

Proof. We take the Laplace transform of (3.8), w.r.t. y , so that we get, by considering (1.9):

$$\frac{\partial}{\partial t} \tilde{f}(\theta, t) = -\lambda \frac{e^{\lambda t}}{\xi} g(\theta) [\tilde{f}(\theta, t)]^2, \quad (3.10)$$

with $\tilde{f}(\theta, 0) = \frac{\xi}{\xi + g(\theta)}$. We can check, by differentiating, that the solution to (3.10) is equal to

$$\tilde{f}(\theta, t) = \frac{\xi e^{-\lambda t}}{\xi e^{-\lambda t} + g(\theta)}. \quad (3.11)$$

On the other hand we can start from the distribution function of (3.9) which can be written as

$$F_{Y_g}(y, t) := P\{Y_g(t) < y\} = \sum_{n=1}^{\infty} P(B(t) = n) F_{X_g}^{*(n)}(y), \quad (3.12)$$

where $F_{X_g}^{*(n)}$ denotes the n -th convolution of $F_{X^{(g)}}(x) := P\{X^{(g)} < x\}$. Under the assumption of absolutely continuous and positive random addends $X_j^{(g)}$, for $j = 1, 2, \dots$, with density $f_{X^{(g)}}(x) := P\{X^{(g)} \in dx\}/dx$, we can write

$$F_{X_g}^{*(n)}(x) = \int_0^{+\infty} F_{X_g}^{*(n-1)}(x-z) f_X(z) dz.$$

By denoting $\tilde{g}(\theta) := \int_0^{+\infty} e^{-\theta x} g(x) dx$ the Laplace transform of $g : \mathbb{R}^+ \rightarrow \mathbb{R}$, we get

$$\tilde{F}_{X_g}^{*(n)}(\theta) = [\tilde{F}_{X_g}(\theta)]^n = \frac{[\tilde{f}_{X_g}(\theta)]^n}{\theta}. \quad (3.13)$$

Therefore we have that

$$\begin{aligned} \tilde{f}_{Y_g}(\theta, t) &= \sum_{n=1}^{\infty} e^{-\lambda t} (1 - e^{-\lambda t})^{n-1} \left[\frac{\xi}{\xi + g(\theta)} \right]^n \\ &= \frac{\xi e^{-\lambda t}}{\xi e^{-\lambda t} + g(\theta)}, \end{aligned} \quad (3.14)$$

which coincides with (3.11). ■

As far as the first-passage time T_β of the process Y_g through the level $\beta > 1$ is concerned, we can derive the general formula of the Laplace transform of its distribution function, by considering (1.5) and by taking into account (3.12) and (3.13), as follows:

$$\begin{aligned}\mathcal{L}[P\{T_\beta < t\}; \theta] &= \mathcal{L}[P\{Y(t) > \beta\}; \theta] = \frac{1}{\theta} - \frac{1}{\theta} \tilde{f}_{Y_g}(\theta, t) \\ &= \frac{g(\theta)}{\theta [e^{-\lambda t} \xi + g(\theta)]}.\end{aligned}\quad (3.15)$$

Remark 3.1 *The following time-changed representation of the process Y_g can be checked, by proving the equality of the one-dimensional distributions:*

$$Y_g(t) \stackrel{d}{=} \mathcal{A}_g(Y(t)), \quad (3.16)$$

for any $t \geq 0$, where Y is the compound birth process (with exponential jumps) defined in (1.4) and supposed independent of the subordinator \mathcal{A}_g . Indeed, the Laplace transform of $\mathcal{A}_g(Y(t))$ can be written, for any $t \geq 0$, as

$$\begin{aligned}\mathbb{E}e^{-\theta \mathcal{A}_g(Y(t))} &= \int_0^{+\infty} \mathbb{E}e^{-\theta \mathcal{A}_g(z)} f_Y(z, t) dz \\ &= \int_0^{+\infty} e^{-g(\theta)z} f_Y(z, t) dz \\ &= [by (2.3)] \\ &= \frac{\xi e^{-\lambda t}}{g(\theta) + \xi e^{-\lambda t}},\end{aligned}$$

which coincides with (3.14). Thus (3.16) follows from the unicity of the Laplace transform.

3.1 Some special cases

(i) The Mittag-Leffler case

Let $g(\theta) = \theta^\alpha$, for $\alpha \in (0, 1]$, then the law of the addends is Mittag-Leffler of parameters α, ξ and, in this special case, we can obtain an explicit and simple formula for the density of $Y_g := Y_\alpha$. Recall that the Mittag-Leffler function, with two parameters, is defined as

$$E_{\beta, \gamma}(x) = \sum_{j=0}^{\infty} \frac{x^{\beta j}}{\Gamma(\beta j + \gamma)}, \quad x, \beta, \gamma \in \mathbb{C}, \quad \operatorname{Re}(\beta), \operatorname{Re}(\gamma) > 0. \quad (3.17)$$

Then the density of X^g can be obtained by applying the well-known formula of the Laplace transform of the Mittag-Leffler function (see [12], formula (1.9.13), for $\rho = 1$), i.e.

$$\mathcal{L}\left\{x^{\gamma-1} E_{\beta, \gamma}(Ax^\beta); s\right\} = \frac{s^{\beta-\gamma}}{s^\beta - A}, \quad (3.18)$$

with $\operatorname{Re}(\beta), \operatorname{Re}(\gamma) > 0$, $A \in \mathbb{R}$ and $s > |A|^{1/\operatorname{Re}(\beta)}$. Indeed we have

$$f_{X^{(\alpha)}}(x) = \mathcal{L}^{-1}\left\{\frac{\xi}{\xi + \theta^\alpha}; x\right\} = \xi x^{\alpha-1} E_{\alpha, \alpha}(-\xi x^\alpha), \quad x, \xi > 0, \quad \alpha \in (0, 1], \quad (3.19)$$

which is the density of $\mathcal{A}_\alpha(X)$, where \mathcal{A}_α is the α -stable subordinator and X an independent exponential r.v. The equation (3.8), in this case, reduces to the following space-fractional differential equation of order α ,

$$\frac{\partial}{\partial t} f_{Y_\alpha}(y, t) = -\lambda \frac{\partial^\alpha}{\partial y^\alpha} [f_{Y_\alpha} * f_{Y_\alpha}](y, t), \quad y, t > 0, \quad (3.20)$$

with initial condition $f_{Y_\alpha}(y, 0) = \xi x^{\alpha-1} E_{\alpha, \alpha}(-\xi y^\alpha)$.

The derivative appearing in (3.20) is the Caputo fractional derivative defined as follows: let $\alpha > 0$, $m = \lfloor \alpha \rfloor + 1$ and assume that $u : [a, b] \rightarrow \mathbb{R}$, $b > a$, is an absolutely continuous function, with absolutely continuous derivatives up to order m on $[a, b]$, then, for $x \in [a, b]$,

$$\frac{d^\alpha}{dx^\alpha} u(x) := \begin{cases} \frac{1}{\Gamma(m-\alpha)} \int_a^x \frac{1}{(x-s)^{\alpha-m+1}} \frac{d^m}{ds^m} u(s) ds, & \alpha \notin \mathbb{N}_0 \\ \frac{d^m}{dx^m}, & \alpha = m \in \mathbb{N}_0 \end{cases} \quad (3.21)$$

is the Caputo fractional derivative of order α (see [12], p.92). Indeed, for $g(\theta) = \theta^\alpha$, we have that

$$\mathcal{D}_x^g = \frac{d^\alpha}{dx^\alpha}, \quad (3.22)$$

as it can be checked by considering (1.8) with the Lévy measure $\bar{\nu}(ds) = \alpha s^{-\alpha-1} ds / \Gamma(1-\alpha)$ and the tail Lévy measure $\nu(ds) = s^{-\alpha} ds / \Gamma(1-\alpha)$ (see Remark 2.6 in [24] for details).

The density of the process (3.9), which we denote now as Y_α , is given by

$$f_{Y_\alpha}(y, t) = \xi e^{-\lambda t} y^{\alpha-1} E_{\alpha, \alpha}(-\xi e^{-\lambda t} y^\alpha), \quad y, t \geq 0, \quad \alpha \in (0, 1], \quad (3.23)$$

as can be obtained by inverting the Laplace transform (3.14), by means of (3.18), with $g(\theta) = \theta^\alpha$ and $\beta = \gamma = \alpha$.

The Mittag-Leffler r.v. has infinite moments and thus the same holds for the process Y_α . This is confirmed by the representation $\mathcal{A}_\alpha(Z)$ of the random addends, since it is well-known that the stable law has infinite moments of order greater than α .

The distribution of the first-passage time through the level $\beta > 1$ in this case can be obtained, by considering (3.15), which reduces to

$$\mathcal{L}[P\{T_\beta < t\}; \theta] = \frac{\theta^{\alpha-1}}{\theta^\alpha + \xi e^{-\lambda t}}. \quad (3.24)$$

By inverting (3.24) we get

$$P\{T_\beta < t\} = \begin{cases} 0, & t \leq 0 \\ E_{\alpha, 1}(-\xi e^{-\lambda t} \beta^\alpha), & t > 0 \end{cases}, \quad (3.25)$$

which coincides with (2.5) for $\alpha = 1$. The jump in the origin of (3.25) coincides with $P\{T_\beta = 0\} = E_{\alpha, 1}(-\xi \beta^\alpha) = P(X^{(\alpha)} > \beta)$.

Moreover, the previous probability converges to zero, as β tends to infinity, but with a power law (instead of exponentially), as can be checked by recalling the well-known asymptotic behavior of the Mittag-Leffler function (see [12], formula (1.8.11)), for $|z| \rightarrow +\infty$, i.e.

$$E_{\beta, \gamma}(z) = -\sum_{k=1}^n \frac{z^{-k}}{\Gamma(\gamma - \beta k)} + O(z^{-n-1}), \quad n \in \mathbb{N}, \quad (3.26)$$

where $0 < \beta < 2$, $\mu \leq \arg(z) \leq \pi$, $\pi\beta/2 < \mu < \min\{\pi, \pi\beta\}$. Thus, for $\beta \rightarrow +\infty$, we get $P\{T_\beta < t\} \simeq O(\beta^{-\alpha})$.

The density of the absolutely continuous component is easily obtained by taking the first derivative of (3.25) and reads

$$f_{T_\beta}(t) = \frac{\lambda \xi e^{-\lambda t} \beta^\alpha}{\alpha} E_{\alpha,\alpha}(-\xi e^{-\lambda t} \beta^\alpha), \quad t > 0.$$

We thus obtain the definition of the following fractional extension of the reflected Gumbel density

$$f(x) = \frac{e^{-x}}{\alpha E_{\alpha,\alpha+1}(-\beta^\alpha)} E_{\alpha,\alpha}(-e^{-x} \beta^\alpha), \quad x > 0, \quad (3.27)$$

where the normalizing constant is obtained as follows

$$\int_0^{+\infty} e^{-x} E_{\alpha,\alpha}(-e^{-x} \beta^\alpha) dx = \int_0^1 E_{\alpha,\alpha}(-z \beta^\alpha) dz = \alpha E_{\alpha,\alpha+1}(-\beta^\alpha).$$

The expected first-passage time through β can be obtained from (3.25), as follows:

$$\begin{aligned} \mathbb{E}T_\beta &= - \int_0^{+\infty} \sum_{j=1}^{\infty} \frac{(-\xi e^{-\lambda t} \beta^\alpha)^j}{\Gamma(\alpha j + 1)} dt = \frac{\xi \beta^\alpha}{\lambda} \sum_{l=0}^{\infty} \frac{(-\xi \beta^\alpha)^l (\Gamma(l+1))^2}{l! \Gamma(\alpha l + \alpha + 1) \Gamma(l+2)} \\ &= \frac{\xi \beta^\alpha}{\lambda} {}_2\Psi_2 \left[-\xi \beta^\alpha \mid \begin{matrix} (1, 1) & (1, 1) \\ (\alpha + 1, \alpha) & (2, 1) \end{matrix} \right] \end{aligned}$$

which reduces to (2.6) for $\alpha = 1$. Its asymptotic behavior can be obtained by applying Theorem 1.2, p.19 in [17], for $\mu = \alpha > 0$ and $d = \min\{-1, -1\} = -1$ and $\alpha > 0$, we get, for $\beta \rightarrow +\infty$,

$$\begin{aligned} \mathbb{E}T_\beta &= \frac{\xi \beta^\alpha}{\lambda} H_{2,3}^{1,2} \left[\xi \beta^\alpha \mid \begin{matrix} (0, 1) & (0, 1) \\ (0, 1) & (-\alpha, \alpha) & (-1, 1) \end{matrix} \right] \\ &\simeq \frac{\xi \beta^\alpha}{\lambda} O((\beta^\alpha)^{-1}) = \frac{\xi}{\lambda} O(1). \end{aligned}$$

Thus the expected first-passage time through the level β converges to the same limit of the exponential case, for any α , even though the expected value of Y_α is infinite, while it is finite for $\alpha = 1$.

(ii) The tempered case

We consider now the case $g(\theta) = (\mu + \theta)^\alpha - \mu^\alpha$, for $\mu > 0$, which is the Bernstein function of the tempered α -stable subordinator $\mathcal{A}_{\alpha,\mu}$, for $\alpha \in (0, 1]$. The law of the addends can be written explicitly, by the well-known relationship between the density of the tempered stable $h_{\alpha,\mu}(x, t)$ and that of the stable itself, i.e.

$$h_{\alpha,\mu}(x, t) = \exp\{-\mu x - \mu^\alpha t\} h_\alpha(x, t), \quad x, t \geq 0.$$

Since, in this case,

$$Y_g(0) \stackrel{d}{=} \mathcal{A}_{\alpha,\mu}(X),$$

where X is again exponentially distributed with parameter ξ and independent of $\mathcal{A}_{\alpha,\mu}$, we have that

$$f_{X_\mu^{(\alpha)}}(x) = \xi e^{-\mu x} \int_0^{+\infty} e^{-\xi t - \mu^\alpha t} h_\alpha(x, t) dt = \xi e^{-\mu x} x^{\alpha-1} E_{\alpha,\alpha}(-(\xi - \mu^\alpha)x^\alpha), \quad (3.28)$$

which generalizes (3.19), for $\mu \neq 1$. Analogously to the previous case, we can write the transition density of the process $Y_g := Y_{\alpha,\mu}$ as

$$f_{Y_{\alpha,\mu}}(y, t) = \xi e^{-\lambda t - \mu y} y^{\alpha-1} E_{\alpha,\alpha}(-(\xi e^{-\lambda t} - \mu^\alpha)y^\alpha), \quad y, t \geq 0, \quad \alpha \in (0, 1]. \quad (3.29)$$

Moreover, from Theorem 3.2, we get that (3.29) satisfies equation (3.20), where the fractional α -order Caputo derivative must be replaced by the following Caputo-type tempered derivative

$$\frac{d^{\alpha,\mu}}{dx^{\alpha,\mu}} u(x) := \frac{\alpha \mu^\alpha}{\Gamma(1-\alpha)} \int_0^x \Gamma(-\alpha; \mu s) \frac{d}{ds} u(s) ds, \quad \alpha \in (0, 1), \quad \mu > 0,$$

(where $\Gamma(\eta, x) := \int_x^{+\infty} e^{-t} t^{\eta-1} dt$ is the upper incomplete Gamma function). Indeed the tail Lévy measure reads $\nu(ds) = \frac{\alpha \mu^\alpha \Gamma(-\alpha; \mu s) ds}{\Gamma(1-\alpha)}$ (see [24]). In this case, the Laplace transform of the first-passage time distribution (3.15) reduces to

$$\mathcal{L}[P\{T_\beta < t\}; \theta] = \frac{(\mu + \theta)^\alpha - \mu^\alpha}{\theta [e^{-\lambda t} \xi + (\mu + \theta)^\alpha - \mu^\alpha]}. \quad (3.30)$$

By inverting (3.30) and denoting by $\gamma(a; x) := \int_0^x e^{-t} t^{a-1} dt$ the lower incomplete Gamma function, we can write that, for $t > 0$,

$$\begin{aligned} P\{T_\beta < t\} &= 1 - \frac{\xi e^{-\lambda t}}{\mu^\alpha} \sum_{j=0}^{\infty} \left(-\frac{\xi e^{-\lambda t} - \mu^\alpha}{\mu^\alpha} \right)^j \frac{\gamma(\alpha j + \alpha; \mu \beta)}{\Gamma(\alpha j + \alpha)} \\ &= 1 - \frac{\xi \beta^\alpha e^{-\lambda t}}{\mu^\alpha} \sum_{j=0}^{\infty} \left(-\frac{\beta^\alpha (\xi e^{-\lambda t} - \mu^\alpha)}{\mu^\alpha} \right)^j E_{1,\alpha j + \alpha + 1}(\mu \beta). \end{aligned}$$

where we have considered formula (3.7) p.316 in [18] together with formula (4.2.8) in [10]. We now compare (in Fig.2) the pdf's of the compound birth process in the two special cases of Mittag-Leffler and tempered stable addends (given in (3.23) and (3.29), respectively) with that of the exponential case: with respect to the latter, the pdf's fall generally quicker while they have longer tails. Moreover the presence of the tempering parameter in the tempered stable case causes both an initial slower fall of the density compared to the pure α -stable case and faster fall for large values of y .

In Fig.3 we explore the influence of the tempering parameter μ for the pdf given in (3.29), in the tempered case: as expected, the greater the value of μ the faster the fall of the tails. Finally, we compare the first crossing time probability of Y through the level β , in the usual three cases, i.e. exponential, stable and tempered stable, by plotting it with respect to time (in Fig.4) and to β (in Fig.5).




Figure 2: The pdf's of the compound birth process Y , with exponential, Mittag-Leffler and tempered stable addends: for $t = 1, \lambda = 1, \xi = 0.5, \alpha = 0.2(a), 0.5(b), 0.8(c), \mu = 10$.




Figure 3: The pdf's of the compound birth process Y_g , with tempered stable addends, for different values of μ and for $\xi = \lambda = 1, \alpha = 0.8, t = 1$.




Figure 4: The first crossing probability of the level β , for $\mu = 5, \xi = \lambda = 1, \alpha = 0.5, t = 1$.




Figure 5: The first crossing probability of the level $\beta=2$, for $\mu = 5, \xi = \lambda = 1, \alpha = 0.5$.

(iii) The gamma case

For $g(\theta) = \log\left(1 + \frac{\theta}{b}\right)$, for $b > 0$, which is the Bernstein function associated to the gamma distribution, the law of the addends X_Γ can be written as follows:

$$f_{X_\Gamma}(x) = \xi e^{-bx} \int_0^{+\infty} \frac{b^t}{\Gamma(t)} x^{t-1} e^{-\xi t} dt$$

with mean value $\mathbb{E}X_\Gamma = 1/\xi b$ and Laplace transform

$$\begin{aligned} \tilde{f}_{X_\Gamma}(\theta) &= \xi \int_0^{+\infty} e^{-bx-\theta x} \int_0^{+\infty} \frac{b^t}{\Gamma(t)} x^{t-1} e^{-\xi t} dt dx \\ &= \xi \int_0^{+\infty} \left(\frac{be^{-\xi}}{b+\theta}\right)^t dt = \frac{\xi}{\log(b+\theta) - \log(be^{-\xi})}. \end{aligned} \quad (3.31)$$

In this case, by considering that

$$\log\left(1 + \frac{\theta}{b}\right) = \int_0^{+\infty} (1 - e^{-\theta x}) x^{-1} e^{-bx} dx,$$

we can write the tail Lévy measure as

$$\nu(ds) = ds \int_s^{+\infty} z^{-1} e^{-bz} dz = ds \int_{bs}^{+\infty} \omega^{-1} e^{-\omega} d\omega = E_1(bs) ds,$$

where $E_1(x) = -\text{Ei}(-x)$ and $\text{Ei}(x)$ denotes the exponential integral. Thus the differential equation governing the process (1.4) coincides with (2.2) with convolution-type derivative defined as follows:

$$\mathcal{D}_t^g u(t) := \int_0^t \frac{d}{ds} u(t-s) E_1(bs) ds.$$

In this case the Laplace transform of the first-passage time distribution (3.15) reduces to

$$\mathcal{L}[P\{T_\beta < t\}; \theta] = \frac{\log(\theta + b) - \log b}{\theta [e^{-\lambda t} \xi + \log(\theta + b) - \log b]},$$

which can be inverted by considering (3.31), as follows, for $t > 0$:

$$\begin{aligned} P\{T_\beta < t\} &= \xi e^{-\lambda t} \int_\beta^{+\infty} e^{-by} \int_0^{+\infty} \frac{b^z}{\Gamma(z)} y^{z-1} \exp\{-\xi e^{-\lambda t} z\} dz dy \\ &= \xi e^{-\lambda t} \int_0^{+\infty} \frac{\Gamma(z; \beta b)}{\Gamma(z)} \exp\{-\xi e^{-\lambda t} z\} dz. \end{aligned}$$

The expected first-passage time through β can be written as follows:

$$\begin{aligned} \mathbb{E}T_\beta &= \xi \int_0^{+\infty} \frac{\gamma(z; \beta b)}{\Gamma(z)} \int_0^{+\infty} e^{-\lambda t} \exp\{-\xi e^{-\lambda t} z\} dt dz \\ &= \frac{\xi}{\lambda} \int_0^{+\infty} \frac{\gamma(z; \beta b)}{\Gamma(z)} \int_0^1 \exp\{-\xi uz\} du dz \\ &= \frac{1}{\lambda} \int_0^{+\infty} \frac{\gamma(z; \beta b)}{\Gamma(z+1)} (1 - e^{-\xi z}) dz. \end{aligned}$$

By applying the monotone convergence theorem and considering that $\frac{\gamma(z; \beta b)}{\Gamma(z+1)} \rightarrow \frac{1}{z}$, as $\beta \rightarrow \infty$, it is easy to check that $\mathbb{E}T_\beta$ is infinite in the limit.

(iv) The Poisson case

Let $g(\theta) = \lambda(1 - e^{-\theta})$, for $\lambda > 0$, which is the Bernstein function associated to the Poisson distribution and with Lévy measure $\nu(s) = \lambda\delta(s-1)$, where $\delta(\cdot)$ is the Dirac's delta function. In this case, since the distribution of the addends $X^{(\lambda)}$ is discrete and integer valued, we must adapt the notation of the previous sections: let $p_x^{(\lambda)} := P\{X^{(\lambda)} = x\}$ denote the addends' probability mass function and let $\tilde{p}_X^{(\lambda)}(\theta) := \sum_{x=0}^{\infty} e^{-\theta x} p_x^{(\lambda)}$, then (3.13) must be replaced by

$$\tilde{F}_X^{*(n)}(\theta) = \frac{[\tilde{p}_X^{(\lambda)}(\theta)]^n}{\theta}.$$

Moreover, formula (3.7) is substituted by

$$p_x^{(\lambda)} = \xi \int_0^{+\infty} e^{-\xi t} P\{N(t) = x\} dt$$

Correspondingly, we have, for $Y_\lambda(t) = \sum_{j=1}^{B(t)} X_j^{(\lambda)}$, that $q_x^{(\lambda)}(t) := P\{Y_\lambda(t) = x\}$ with Laplace transform

$$\tilde{q}_Y^{(\lambda)}(\theta) := \sum_{x=0}^{\infty} e^{-\theta x} q_x^{(\lambda)}(t) = \frac{\xi e^{-\lambda t}}{\xi e^{-\lambda t} + g(\theta)} = \frac{\xi e^{-\lambda t}}{\xi e^{-\lambda t} + \lambda(1 - e^{-\theta})}. \quad (3.32)$$

It is easy to see that, in this special case, the addends follow a geometric distribution of parameter $p = \xi/(\xi + \lambda)$: indeed, here $X^{(\lambda)} \stackrel{d}{=} N(X)$, with $X \sim \text{Exp}(\xi)$ independent of N , so that the probability mass function of $X^{(\lambda)}$ can be written as

$$p_x^{(\lambda)} = \xi \int_0^{+\infty} e^{-\xi t} P\{N(t) = x\} dt = \frac{\xi}{\lambda + \xi} \left(\frac{\lambda}{\lambda + \xi} \right)^x, \quad x = 0, 1, \dots \quad (3.33)$$

The distribution of the process Y_λ can be written, for $y = 0, 1, \dots$ and for any $t \geq 0$, as

$$q_y^{(\lambda)}(t) = \frac{\xi e^{-\lambda t} \lambda^y}{y!} \int_0^{+\infty} e^{-(\xi e^{-\lambda t} + \lambda)z} z^y dz = \frac{\xi e^{-\lambda t} \lambda^y}{(\xi e^{-\lambda t} + \lambda)^{y+1}}, \quad (3.34)$$

which coincides with a geometric probability mass function with parameter $p = \xi e^{-\lambda t} / (\xi e^{-\lambda t} + \lambda)$. This can be checked by evaluating the Laplace transform of (3.34), which coincides with (3.32).

The distribution function of the first-passage time through the level β is given by

$$P\{T_\beta < t\} = \begin{cases} 0, & t \leq 0 \\ \left(\frac{\lambda}{\lambda + \xi e^{-\lambda t}}\right)^{\beta+1}, & t > 0 \end{cases},$$

which converges to zero, for $\beta \rightarrow +\infty$.

As far as the differential equation satisfied by the survival function of $N(X)$, we can specialize the definition (1.10), by considering that $\nu(ds) = \lambda ds \int_s^{+\infty} \delta(x-1) dx = \lambda ds 1_{(-\infty, 1]}(s)$ and thus

$$\mathcal{D}_t^g u(t) = \lambda \int_0^{t \wedge 1} \frac{d}{ds} u(t-s) ds = \lambda \begin{cases} u(t) - u(0), & t < 1 \\ u(t) - u(t-1), & t \geq 1 \end{cases}$$

Its Laplace transform reads

$$\begin{aligned} & \int_0^{+\infty} e^{-\theta t} \lambda [u(t) - u((t-1) \vee 0)] dt & (3.35) \\ &= \lambda \tilde{u}(\theta) - \lambda \int_0^1 e^{-\theta t} u(0) dt - \lambda \int_1^{+\infty} e^{-\theta t} u(t-1) dt \\ &= \lambda(1 - e^{-\theta}) \tilde{u}(\theta) - \frac{\lambda(1 - e^{-\theta})}{\theta} u(0), \end{aligned}$$

which coincides with (1.9), for this choice of g . As a consequence, by taking into account Lemma 3.1, we obtain that the survival function of the addends $P\{X^{(\lambda)} \geq x\}$ satisfies the following equation (for $x = 1, 2, \dots$)

$$(\lambda + \xi)u(x) = \lambda u(x-1)$$

with $u(0) = 1$, as can be checked also directly by considering that $P\{X^{(\lambda)} \geq x\} = \left(\frac{\lambda}{\lambda + \xi}\right)^x$, in this case. Theorem 3.2 can be formulated as follows, with the convention that $q_{-1}(t) = 0$: the solution to the initial value problem

$$\begin{aligned} \frac{\partial}{\partial t} q_y(t) &= -\lambda^2 \frac{e^{\lambda t}}{\xi} [(q_y(t) * q_y(t)) - (q_{y-1}(t) * q_{y-1}(t))] & (3.36) \\ &= -\lambda^2 \frac{e^{\lambda t}}{\xi} [I - \Delta] (q_y(t) * q_y(t)) \end{aligned}$$

(with $t \geq 0$, $y = 0, 1, \dots$, and $q_y(0) = \xi \lambda^y / (\xi + \lambda)^{y+1}$), coincides with (3.34). This can be checked either directly, by considering that $q_y^{(\lambda)}(t) * q_y^{(\lambda)}(t) = \xi^2 e^{-2\lambda t} \lambda^y y / (\xi e^{-\lambda t} + \lambda)^{y+2}$, or by taking the Laplace transform and verifying that the discrete analogue of (3.10) holds in this case, i.e.

$$\frac{\partial}{\partial t} \tilde{q}_Y(\theta) = -\lambda^2 \frac{e^{\lambda t}}{\xi} (1 - e^{-\theta}) [\tilde{q}_Y(\theta)]^2, \quad (3.37)$$

for any $t \geq 0$.

References

- [1] Beghin L., Fractional relaxation equations and Brownian crossing probabilities of a random boundary, *Adv. Appl. Probab.*, 2012, 44, 479- 505.
- [2] Beghin, L., Macci, C., Alternative forms of compound fractional Poisson processes, *Abstr. Appl. Anal.*, 2012, ID 747503, 30 p.
- [3] Buchak K., Sakhno L., Compositions of Poisson and gamma processes, *Modern Stochastics: Theory and Applications*, 2017, 4 (2), 161-188.
- [4] Cahoy D.O., Polito F., Simulation and estimation for the fractional Yule process, *Methodology and Computing in Applied Probability*, 2012, 14 (2), 383-403.
- [5] Cahoy D.O., Uchaikin V.V., Woyczynski W.A., Parameter estimation for fractional Poisson processes, *J. Statist. Plann. Inference*, 2010, 140(11), 3106-3120.
- [6] Crowder M., *Multivariate Survival Analysis and Competing Risks*, 2012, Taylor and Francis.
- [7] Devroye L., Random variate generation for exponentially and polynomially tilted stable distributions, *ACM Trans. Model. Comput. Simul.*, 2009, 19(4), 1-20.
- [8] Gajda J., Wyłomańska A., Time-changed Ornstein-Uhlenbeck process, *J. Phys. A: Math. Theor.*, 2015, 48, 1-19.
- [9] Glockle W.G., Nonnenmacher T.F., Fox function representation of non-Debye relaxation processes, *J Stat Phys.*, 1993, 71 (3/4), 741-757.
- [10] Gorenflo R. , Kilbas A.A., Mainardi F., Rogosin S.V., *Mittag-Leffler Functions, Related Topics and Applications*, 2014, Springer-Verlag, Berlin Heidelberg.
- [11] Hofert M., Sampling exponentially tilted stable distributions, *ACM Trans. Model. Comput. Simul.*, 2011, 22(1) 3-11.
- [12] Kilbas A.A., Srivastava H.M., Trujillo J.J., *Theory and Applications of Fractional Differential Equations*, vol. 204 of North-Holland Mathematics Studies, 2006, Elsevier Science B.V., Amsterdam.
- [13] Kochubei A.N., General fractional calculus, evolution equations and renewal processes, *Integral Equations Operator Theory*, 2011,71, 583-600.
- [14] Lansky P., Polito F., Sacerdote L., Generalized nonlinear Yule models, *J Stat Phys.*, 2016, 165, 661–679.
- [15] Maheshwari A., Vellaisamy P., Fractional Poisson process time-changed by Lévy subordinator and its inverse, *Journal of Theoretical Probability*, 2019, 32(3), 1278-1305.
- [16] Mainardi F., Mura A., Gorenflo R., Stojanović M., The two forms of fractional relaxation of distributed order, *Journal of Vibration and Control*, 2007, 13, (9-10), 1249-1268.

- [17] Mathai A.M., Saxena R.K., Haubold H.J., *The H-Function: Theory and Applications*, 2010, Springer, New York.
- [18] Miller K.S., Ross B., *An Introduction to the Fractional Calculus and Fractional Differential Equations*, 1993, John Wiley and Sons, New York.
- [19] Orsingher E., Polito F., Fractional pure birth process, *Bernoulli*, 2010, 16, (3), 858-881.
- [20] Orsingher E., Polito F., On a fractional linear birth-death process. *Bernoulli*, 2011, 17, (1), 114–137.
- [21] Orsingher E., Ricciuti C., Toaldo B., On semi-Markov processes and their Kolmogorov’s integro-differential equations, *Journ. Funct. Analysis*, 2018, 275 (4), 830-868.
- [22] Sendova K.P., Minkova L.D., Introducing the nonhomogeneous compound-birth process, *Stochastics*, 2019, 1-19.
- [23] Sobczyk K., Spencer, J., *Random Fatigue: From Data to Theory*, 1992, Academic Press, New York.
- [24] Toaldo B., Convolution-type derivatives, hitting-times of subordinators and time-changed C_0 -semigroups, *Potential Analysis*, 2015, 42, 1, 115-140.